

Crossover from Conserving to Lossy Transport in Circular Random Matrix Ensembles

Steven H. Simon^a and Aris L. Moustakas^b

^a*Lucent Technologies, Bell Labs, Murray Hill, NJ, 07974, USA*

^b*Department of Physics, University of Athens, Panepistimiopolis, Athens 15784, Greece*

In a quantum dot with three leads the transmission matrix $\mathbf{t}_{12}$ between two of these leads is a truncation of a unitary scattering matrix $\mathbf{S}$, which we treat as random. As the number of channels in the third lead is increased, the constraints from the symmetry of $\mathbf{S}$ become less stringent and $\mathbf{t}_{12}$ becomes closer to a matrix of complex Gaussian random numbers with no constraints. We consider the distribution of the singular values of $\mathbf{t}_{12}$, which is related to a number of physical quantities.

Random matrix theory has enjoyed enormous success in a wide range of fields of physics[1]. One impressive success has been in mesoscopic physics[2], where the scattering matrix between two leads (a source and a sink) of a disordered quantum dot has been modelled as a random matrix. Such scattering matrices can be from the circular unitary ensemble (CUE or unitary matrices) if time reversal symmetry is broken, the circular orthogonal ensemble (COE or unitary symmetric matrices) if time reversal is unbroken, or the circular symplectic ensemble (CSE or unitary self dual matrices) if spin-orbit scattering is present and time reversal is unbroken[2]. Another success of random matrix theory has been in the arena of radio communications[3, 4] where transmission matrices between transmitting and receiving antennas have been modelled as random matrices with all matrix elements being independent and identically distributed (*i.i.d.*) complex Gaussian. These two examples are, in fact, quite closely related to each other in that both stem from wave transmission problems. The fundamental difference between these two cases is that the former is “conserving” while the latter is “lossy”. When an electron enters a dot, it must exit one of the two leads whereas in the radio communication problem most of the photons which are transmitted are lost to the environment and neither reach the receiver nor return to the transmitter. In this manuscript we will study the crossover between these two cases by studying a quantum dot with a “third lead” grounded at the same voltage as the sink of the dot. As the number of channels to ground through the third lead is increased (i.e., the amount of loss is increased), the statistics of transmission between the source and the sink become more and more similar to the *i.i.d.* complex Gaussian case *independent* of the underlying symmetry (COE, CUE, or CSE). We note that this application of a “third lead” is quite distinct from that used previously in the mesoscopic literature to model dephasing[2]. In that case, a constraint is imposed that no current may flow out the third lead; whereas in the present case, we allow current to flow out the third lead, giving us true loss of electrons. We will not consider dephasing here, so this work strictly applies to dots at zero temperature.

We thus consider a quantum dot with three leads, a source (lead 1) at voltage V , a grounded sink (lead 2)

at voltage 0, and a loss lead (lead 3) also grounded at voltage 0. We assume there are N_i fully open channels going in (and out of) lead i . It is useful to define

$$N = N_1 + N_2 + N_3, \quad (1)$$

$$N_m = \min(N_1, N_2), \quad N_\Delta = |N_1 - N_2|. \quad (2)$$

The $N \times N$ scattering matrix between all three leads can be written in block form as

$$\mathbf{S} = \begin{pmatrix} \mathbf{r}_{11} & \mathbf{t}_{12} & \mathbf{t}_{13} \\ \mathbf{t}_{21} & \mathbf{r}_{22} & \mathbf{t}_{23} \\ \mathbf{t}_{31} & \mathbf{t}_{32} & \mathbf{r}_{33} \end{pmatrix} \quad (3)$$

where $\mathbf{r}_{ii}$ is $N_i \times N_i$ and $\mathbf{t}_{ij}$ is $N_i \times N_j$. A quantity of interest is the $N_1 \times N_2$ matrix $\mathbf{t}_{12}$. Making a singular value decomposition (SVD) we have $\mathbf{t}_{12} = \mathbf{U}\mathbf{D}\mathbf{V}$ where $\mathbf{U}$ is an N_1 dimensional unitary matrix, $\mathbf{V}$ is an N_2 dimensional unitary matrix, and $\mathbf{D}$ is an $N_1 \times N_2$ matrix with the (real) singular values $d_i > 0$ along the N_m diagonal elements and zeros elsewhere. More physically relevant perhaps is the Hermitian matrix

$$\mathbf{T}_{12} = \mathbf{t}_{12}^\dagger \mathbf{t}_{12} = \mathbf{V}^\dagger \mathbf{D}^\dagger \mathbf{D} \mathbf{V} \quad (4)$$

whose N_m nonzero eigenvalues $z_i = d_i^2$ correspond to the squares of the singular values of $\mathbf{t}_{12}$. It is convenient to work in terms of the eigenvalues $\{z_i\}$ rather than the singular values $\{d_i\}$. Many physical quantities are given in terms of these eigenvalues[2, 5]. For example, the current out of lead 2 is given by $I_2 = g_{12}G_0V$ where $G_0 = 2e^2/h$ and

$$g_{12} = \text{Tr}[\mathbf{T}_{12}] = \sum_{i=1}^{N_m} z_i \quad (5)$$

Similarly the shot noise of the current at lead 2 can be shown[2, 5] to be given by $2eVG_0s$ with $s = \text{Tr}[\mathbf{T}_{12}(\mathbf{1} - \mathbf{T}_{12})] = \sum_{i=1}^{N_m} z_i(1 - z_i)$. These physical quantities are examples of so-called “linear statistics”[2] whose expectations can be written as an integral over the eigenvalue density $\rho(z)$ of the nonzero eigenvalues of $\mathbf{T}_{12}$. Other linear statistics can also be shown to give physically interesting quantities related to higher moments of the current[2, 5, 6].

We note that a typical element of the matrix $\mathbf{S}$ has magnitude $1/N^{1/2}$ (since $\mathbf{S}\mathbf{S}^\dagger = \mathbf{1}$), so increasing N_3

has the effect of decreasing all the elements of $\mathbf{t}_{12}$. In addition to this normalization change, qualitatively we should expect that the symmetry (COE, CUE, or CSE) constraints on $\mathbf{S}$ become less visible in the submatrix $\mathbf{t}_{12}$ as N_3 is increased, since the constraints are more easily satisfied by the large number of additional variables in the other blocks of $\mathbf{S}$. Thus, we expect that $\mathbf{t}_{12}$ should start to look like an *i.i.d.* random complex Gaussian matrix as N_3 becomes large. When $\mathbf{t}_{12}$ is *i.i.d.* random complex Gaussian, the corresponding $\mathbf{T}_{12}$ is known as a “complex Wishart” matrix[7, 8, 9].

As is frequently the case in random matrix problems, it is easiest to make further progress if one focuses on $\mathbf{S}$ being from the unitary ensemble (broken time reversal invariance). In this case we use a method analogous to that of Ref. [10] to determine the joint probability density (JPD) of the nonzero eigenvalues $\{z_i\}$ of $\mathbf{T}_{12}$. We start by writing the probability density $\mathcal{P}(\mathbf{S}) \propto \delta(\mathbf{S}^\dagger \mathbf{S} - \mathbf{1})$ where we have an N^2 dimensional complex δ function to enforce unitarity. Analogous to [10] we integrate out $\mathbf{r}_{33}, \mathbf{r}_{22}, \mathbf{t}_{31}, \mathbf{t}_{32}, \mathbf{t}_{21}, \mathbf{t}_{23}$ to obtain

$$\mathcal{P}(\mathbf{t}_{12}) \propto \int d\mathbf{t}_{13} \int d\mathbf{r}_{11} \delta(\mathbf{r}_{11}^\dagger \mathbf{r}_{11} + \mathbf{t}_{12}^\dagger \mathbf{t}_{12} + \mathbf{t}_{13}^\dagger \mathbf{t}_{13} - \mathbf{1}) \quad (6)$$

with the delta function being N_1^2 complex dimensional, and the integrals being $N_1(N_1 + N_3)$ complex dimensional. Writing $\mathbf{t}_{12}^\dagger \mathbf{t}_{12}$ in terms of its SVD as in Eq. 4, the Jacobian of this change of variables is given by $J \sim \Delta(\{z\})^2 \prod_i z_i^{N_\Delta}$ where $\Delta(\{z\})$ is the Vandermonde determinant

$$\Delta(\{z\}) = \prod_{1 \leq i < j \leq N_m} (z_i - z_j) = \det[z_i^{j-1}]. \quad (7)$$

We can then integrate out $\mathbf{U}$ and $\mathbf{V}$ to obtain

$$\mathcal{P}(\{z\}) \propto J \int d\mathbf{t}_{13} \int d\mathbf{r}_{11} \delta(\mathbf{r}_{11}^\dagger \mathbf{r}_{11} + \mathbf{t}_{13}^\dagger \mathbf{t}_{12} + \mathbf{z} - \mathbf{1}) \quad (8)$$

where $\mathbf{z}$ is a diagonal matrix of the N_m nonzero eigenvalues z_i (with zeros along the remaining diagonals if $N_1 > N_2$). Performing the remaining integrals yields the final result for the JPD

$$\mathcal{P}(\{z\}) \propto \Delta(\{z\})^2 \prod_{i=1}^{N_m} w(z_i) \quad (9)$$

with weight function

$$w(z) = z^{N_\Delta} (1 - z)^{N_3} \Theta(z) \Theta(1 - z) \quad (10)$$

where the step function $\Theta(x)$ is 1 for $x > 0$ and is zero otherwise. In the limit of large N_3 (with N_Δ and N_m fixed) the factor $(1 - z)^{N_3}$ becomes e^{-zN_3} ($\sim e^{-zN}$ since $z(N_1 + N_2)$ becomes small) as expected for the limiting complex Wishart distribution[1, 7, 8, 9].

It is interesting that Eqs. 9, 10 are similar in form to the distribution found for the (complex) *eigenvalues* of

a (square) truncated unitary matrix found by Ref. [10]. Here, in contrast, we are looking at the *singular* values of the truncated unitary matrix ($\mathbf{t}_{12}$). Despite the similarity of these cases, we will see below that the distribution of the singular values of $\mathbf{t}_{12}$ is *very* different from that of the absolute value of its eigenvalues.

The Vandermonde determinant can be recast as a Slater determinant of a set of wavefunctions $\{\phi_i\}$ which are orthonormal with respect to the weight function $w(z)$

$$\phi_i(z) = C_i P_i^{(N_3, N_\Delta)}(2z - 1) \quad (11)$$

for $i = 0 \dots N_m - 1$ where $P_i^{(a,b)}(x)$ is the Jacobi polynomial[11], and $C_i^2 = [(2i + N_3 + N_\Delta + 1)! (i + N_3 + N_\Delta)!] / [(i + N_3)! (i + N_\Delta)!]$. Thus our distribution is a realization of the so-called Jacobi Unitary Ensemble (JUE)[8, 9]. Interestingly this ensemble occurs in a completely different random matrix problem when one considers the eigenvalues of $A(A + B)^{-1}$ where A and B are both complex Wishart matrices[8, 9]. In the case of the quantum dot with two leads ($N_3 = 0$) it has been previously pointed out[9] that the eigenvalue distribution is a special case of the JUE. Here we have shown that all possible JUEs may be realized in three lead dots.

Using standard techniques[1, 9], once we have our JPD in this orthogonal polynomial form, we can write the kernel $K(z_1, z_2)$ as

$$K(z_1, z_2) = \sqrt{w(z_1)w(z_2)} \sum_{i=0}^{N_m-1} \phi_i(z_1) \phi_i(z_2) \quad (12)$$

and this series can be resummed using the Christoffel-Darboux formula as usual[1, 9] if desired. All correlation functions between the eigenvalues $\{z_i\}$ can be expressed analytically in terms of this kernel[1, 9], the simplest example of which is the normalized single eigenvalue density $\rho(z) = K(z, z)/N_m$ which is shown in Fig. 1. For comparison, the orthogonal case is also shown (data from numerical monte-carlo simulations). In both cases, the distribution clearly converges to the complex Wishart form[7, 8, 9] for large N_3 . Similar convergence to the Wishart form is obtained from numerical simulations for the symplectic case (not shown).

We can also ask about the distribution of $g_{12} = \text{Tr}[\mathbf{T}_{12}]$ which, as mentioned above, corresponds to the conductance of our dot. We have

$$P(g_{12}) = \int d\{z_i\} \delta(g_{12} - \sum_i z_i) \mathcal{P}(\{z\}) \quad (13)$$

$$= \int \frac{dk}{2\pi} \int d\{z_i\} e^{-ik(g_{12} - \sum_i z_i)} \mathcal{P}(\{z\}) \quad (14)$$

It is now convenient to rewrite the $\Delta(\{z_i\})$ in $P(\{z\})$ in its determinant form (Eq. 7) and expand this determinant as an antisymmetrized product. We can then perform the integrals to obtain

$$P(g_{12}) = \frac{1}{\det[\mathbf{F}(0)]} \int \frac{dk}{2\pi} e^{-ik g_{12}} \det[\mathbf{F}(k)] \quad (15)$$




FIG. 1: **Top:** Scaled eigenvalue density $\rho(zN)/N$ of $\mathbf{T}_{12} = \mathbf{t}_{12}^\dagger \mathbf{t}_{12}$. **Bottom:** Scaled distribution $P(g_{12}N)/N$ with $g_{12} = \text{Tr}[\mathbf{T}_{12}]$. In all cases data is shown for $N_1 = 3, N_2 = 2$ and $N_3 = 0, 1, 2, 5, 15$ (note $N = N_1 + N_2 + N_3$). The unitary cases are analytic as discussed in the text. The orthogonal cases ($N_3 \neq 0$) are numerical and the curves have been smoothed for clarity. The orthogonal $N_3 = 0$ case is also analytic as given by Refs. [2, 15]. The dashed curves are the Wishart limit[7] corresponding to $N_3 \rightarrow \infty$ and is the same for both unitary and orthogonal cases. The eigenvalue density is only nonzero for $z < N$, and $P(g_{12}N)$ is only nonzero for $g_{12} < N_m = 2$.

where $\mathbf{F}$ is an $N_m \times N_m$ matrix with elements

$$F_{ab}(k) = \int_0^1 dz z^{a+b-2+N_\Delta} (1-z)^{N_3} e^{ikz} \quad (16)$$

The integrals of Eq. 16 although tedious can always be done, resulting in the form

$$F_{ab}(k) = Q_{ab}(1/k) + e^{ik} R_{ab}(1/k) \quad (17)$$

where Q_{ij} and R_{ij} are simple polynomials of degree $a + b - 1 + N_\Delta + N_3$, and $F_{ij}(0)$ remains finite. We then have

$$\det[\mathbf{F}(k)] = \sum_{n=0}^{N_m} D_n(1/k) e^{ikn} \quad (18)$$

where the D_n 's are polynomials of order $N_m(N - N_m)$. The Fourier transform in Eq. 15 can easily be carried out analytically resulting in $P(g)$ being of the form

$$P(g_{12}) = \sum_{n=0}^{N_m} L_n(g_{12}) \Theta(g_{12} - n) \quad (19)$$

where each L_n is an $N_m(N - N_m) - 1$ degree polynomial. The detailed forms of these polynomials can easily be evaluated on a case-by-case basis (trivially with Mathematica). In the limit of large N_3 , the $(1-z)^{N_3}$ factor in the integral in Eq. 16 can be replaced by e^{-zN_3} and the upper limit of the integral can be extended to ∞ . We then have $F_{ab}(k) = (N_\Delta + a + b - 2)!(N_3 - ik)^{(1-a-b-N_\Delta)}$. The determinant is then $\det \mathbf{F}(k) \sim (N_3 - ik)^{-N_1 N_2}$ resulting in $P(g_{12}) \sim g_{12}^{N_1 N_2 - 1} e^{-g_{12} N_3} \approx g_{12}^{N_1 N_2 - 1} e^{-g_{12} N}$, which is known as the Rayleigh distribution — the characteristic distribution for the trace of a complex Wishart matrix.

In Fig. 1, examples of the distribution $P(g_{12})$ are shown. The results in the unitary case are analytic as described above, whereas the orthogonal data has been generated numerically. Convergence to the complex Wishart (Rayleigh) limit is again clear in both cases. Similar convergence to this limiting form is obtained from numerical simulations for the symplectic case (not shown).

As with most random matrix problems, considerable simplification occurs in the limit of large matrices. We will thus consider the limit of $N_1, N_2, N_3 \rightarrow \infty$ while keeping the ratios N_i/N fixed. We proceed by considering Eq. 9 as a classical one-dimensional gas with Boltzmann weight given by $\mathcal{P}(\{z\})$. Thus, we have a one particle potential given by $e^{-V(z)} = w(z)$, and two particle interaction $e^{-W(z_i, z_j)} = (z_i - z_j)^2$. Using standard methods[1, 9, 12], we derive the saddle point equations

$$V'(z_i) = \sum_{j \neq i} \frac{2}{z_i - z_j} \quad (20)$$

We then introduce the Green's function

$$G(z) = \frac{1}{N_m} \sum_i \frac{1}{z - z_i} = \int dz' \frac{\rho(z')}{z - z'} \quad (21)$$

with $\rho(z) = \pi^{-1} \text{Im } G(z + i0^+)$ being the normalized single eigenvalue density. It is easy to verify[1, 9, 12] that to order $1/N$ we have $G(z)^2 = \sum V'(z_i)/(z - z_i)$ which we can solve analytically resulting in the eigenvalue density

$$\rho(z) = \frac{N}{N_m} \frac{\sqrt{(z - z_-)(z_+ - z)}}{2\pi z(1 - z)} \quad (22)$$

for $z_- < z < z_+$ and $\rho(z) = 0$ otherwise, where

$$z_{\pm} = N^{-2} \left(\sqrt{N_1(N - N_2)} \pm \sqrt{N_2(N - N_1)} \right)^2 \quad (23)$$

We have also derived this simple form by using the free-probability[13] method, which we will present in a separate paper[14]. In fact, we also show in that work that this form is correct in the large N limit *independent* of whether the original scattering matrix $\mathbf{S}$ is from the COE, CUE, or CSE. This result should not be surprising, as it is well known[1, 2] that for many quantities of physical




FIG. 2: Eigenvalue density $\rho(z)$ of $\mathbf{T}_{12} = \mathbf{t}_{12}^\dagger \mathbf{t}_{12}$ for $N_1/N = 1/6$, $N_2/N = 3/6$ and $N_3/N = 2/6$ for $N_1 = 1, 2, 3, 5$ ($N = 6, 12, 18, 30$). The unitary case is calculated analytically as described in the text. The orthogonal case is calculated numerically. For $N_1 = 1, 2$ the orthogonal case numerical data has been smoothed for clarity. For $N_1 = 3$ (circles) and $N_1 = 5$ (pluses), smoothing is found to destroy the oscillations, so the raw histogrammed data is shown. The dashed line is the large N limit (Eq. 22) to which both the unitary and orthogonal cases converge.

interest (such as the average conductance, etc), the ensembles differ from each other only at order unity whereas leading terms will be order N . These small differences are known as the weak localization corrections.

In Fig. 2 we show how the eigenvalue distributions converge to the large N limit, holding N_i/N fixed. Analytic results are shown for the unitary case, and numerical results are shown for the orthogonal case. In both cases we see convergence to the limiting form of Eq. 22 as expected. Similar convergence to this limit is seen numerically for the symplectic case (not shown).

As mentioned above, the problem of calculating the eigenvalue distribution of $\mathbf{T}_{12}$, or equivalently the singular value distribution of $\mathbf{t}_{12}$ appears similar to the problem of calculating the complex eigenvalue distribution of a (square) truncated random unitary matrix derived in [10]. To see the difference between these two problems, let us examine the case of $N_1/N = N_2/N = n \leq 1/2$ in the large N limit for which both calculations are applicable. From our results, we find the squared singular values density decreases monotonically to zero at $z = z_+ = 4n(1 - n)$. In the work of [10], however, it is found that the density of the norm squared of the eigenvalue distribution is a function that *increases* monotonically, then drops to zero discontinuously at $z = n$.

One might attempt to guess at the general form of the JPD, analogous to Eqs. 9 and 10, for the COE and CSE. Indeed, in the case of $N_3 = 0$ a simple generalization exists[2, 15] which involves changing the exponent of z

in the weight $w(z)$ and changing the exponent of the Vandermonde determinant in Eq. 9 to $\beta = 1$ for the COE and to $\beta = 4$ for the CSE case. However, for the $N_3 \neq 0$ problem no such simple form can apply, since for large N_3 we always expect that $\mathbf{T}_{12}$ will be Wishart (ie, $\mathbf{t}_{12}$ is simply an *i.i.d.* complex Gaussian matrix) so that there will always be quadratic ($\beta = 2$) eigenvalue repulsion *independent* of whether $\mathbf{S}$ is from the COE, CUE, or CSE. We thus leave the full solution of the JPD for the COE and CSE as an open problem.

The authors are greatly indebted to A. Lamacraft and B. Simons for pointing us to this crossover as an interesting problem to examine.

-
- [1] For a good review of Random Matrices in Physics, see, for example T. Guhr, A. Muller-Groeling, and H. A. Weidenmuller, Phys. Rep. **299** (1998) 189.
 - [2] C. W. J. Beenakker, Rev. Mod. Phys. **69**, 731-808 (1997)
 - [3] A. Tulino and S. Verdu, *Foundations and Trends in Communications and Information Theory* **1**, 2004.
 - [4] A. L. Moustakas et al, Science, **287** 181 (2000); See also M. J. Stephen and G. Cwilich, Phys. Rev Lett. **59** 285 (1987); A. Z. Genack in *Scattering and Localization of Classical Waves in Random Media*, P. Sheng, ed, World Scientific (1990).
 - [5] S. A. van Langen and M. Büttiker Phys. Rev. **B 56**, R1680-R1683 (1997); Th. Martin and R. Landauer, Phys. Rev. **B 45**, 1742 (1992).
 - [6] H. Lee, L. S. Levitov, and A. Yu. Yakovets Phys. Rev. **B 51**, 4079-4083 (1995)
 - [7] J. Wishart, Biometrika 20A (1928) 32. See, for example, R. A. Janik and M. A. Nowak, J. Phys. **A36** (2003) 3629.
 - [8] I. Dumitriu and A. Edelman. J. Math. Phys. **43**, 5830, (2002).
 - [9] P. Forrester, "Log-gases and Random Matrices", unpublished book in progress, available at <http://www.ms.unimelb.edu.au/~matpjf/matpjf.html>
 - [10] K. Życzkowski and H.-J. Sommers J. Phys. A **33** 2045 (2000).
 - [11] M. Abramowitz and I. A. Stegun, "Handbook of Mathematical Functions", Dover (1964).
 - [12] "Random Matrices", available at <http://www-spht.cea.fr/articles/t01/014/publi.ps.gz>
 - [13] D. V. Voiculescu, Inv. Math. **104** 201 (1991); For a nice discussion of this technique see [3]; M. Debbah et al, IEEE Tran. Inf. Theory **49**, 1293 (2003); R. A. Janik et al, Phys. Rev. **E55**, 4100 (1997);
 - [14] A. L. Moustakas and S. H. Simon, to be published.
 - [15] P. A. Mello and H. U. Baranger, Waves in Random Media **9**, 105 (1999); cond-mat/9812225.